

HIGH-POWERED MEDICINE

2nd Ed.



**EVIDENCE BASED RECOMMENDATIONS FOR OPTIMIZING
PHARMACOTHERAPY IN CHRONIC DISEASE**

Alex Poppen
DOCTOR OF PHARMACY

High-Powered Medicine

Landmark Clinical Trial Reviews

~ 2nd Edition ~

** FREE SAMPLE **

Alex Poppen

Doctor of Pharmacy

Copyright © 2023 High-Powered Solutions, LLC

All rights reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise without written permission from the publisher. It is illegal to copy this book, post it to a website, or distribute it by any other means without permission.

Alex Poppen, PharmD asserts the moral right to be identified as the author of this work.

This book was prepared solely by the author. It is not intended to be a replacement for reading the original articles referenced within or serve as a substitute for more complete clinical resources. Each trial review contains summarized information from the original article determined to be clinically relevant as well as an analysis and recommendation from the author. The trials are cited within each review to give due credit to said authors.

This book is intended to be used for educational purposes only. The author has made every effort to ensure the accuracy of the information presented. However, it is ultimately the responsibility of the reader to assess the appropriateness of the recommendations presented after consideration of patient specific factors and after consideration of any new developments in the field of medicine. It is important to note that any recommendations included in this book are based solely on the results of clinical trials and are not a replacement for one's own clinical judgment when making decisions regarding an individual's care. The author disclaims all responsibility for any errors or damage incurred as a consequence of the use of the information contained in this book.

This book is NOT to be used to make clinical decisions (such as the diagnosis, treatment or prevention of any disease or condition) in any way and it is not a replacement for clinical judgment. This book CANNOT and DOES NOT contain medical/health advice. Any information or recommendations provided by this book are NOT a substitute for professional advice. Accordingly, before taking any actions based upon such information, the author encourages you to consult with the appropriate professionals.

UNDER NO CIRCUMSTANCE SHALL THE AUTHOR HAVE ANY LIABILITY TO YOU FOR ANY LOSS, DAMAGE OR HARM OF ANY KIND INCURRED AS A RESULT OF THE USE OF THIS BOOK OR RELIANCE ON ANY INFORMATION PROVIDED. YOUR USE OF THIS BOOK AND YOUR RELIANCE ON ANY INFORMATION PROVIDED BY THIS BOOK IS SOLELY AT YOUR OWN RISK.

The author has no financial disclosures or conflicts of interest to note at this time.

“Good doctors use both individual clinical expertise and the best available external evidence, and neither alone is enough.”

-David Sackett, MD

Table of Contents

Quick Review of Biostatistics.....	1
4S.....	4
ACCORD-BG.....	6
ALLHAT.....	8
DAPA-HF.....	11
PARADIGM-HF.....	13
PRECISION.....	15
RACE II.....	17
ROCKET AF.....	19
SPRINT.....	22
VAST-D.....	25

Quick Review of Biostatistics

Please note - this is not meant to be a comprehensive review of biostatistics. It is only meant to serve as a quick reference relevant to this book.

Throughout this book I will be using “significant” in place of “statistically significant” whenever possible in discussing trial results. This was in effort to avoid the redundancy of specifying the *statistical* significance of *statistical* results. Furthermore, **statistical significance does not equal clinical significance** and I did not want to place emphasis on statistical results alone.

Power: the ability of a *trial* to detect a statistically significant difference between groups when a difference truly exists.

- A trial that sets power at 90% (and meets said criteria) means that there is a 90% probability that the observed difference (or lack thereof) between two groups is not due to chance
- It also means there is a 10% chance for showing no difference between groups when a difference truly exists (false negative)
- If power is set but not met and there is no significant difference between treatment groups the results should be considered inconclusive because the trial did not have enough power to confirm that the lack of difference was not due to chance alone
- If power is not met but a statistically significant difference is observed, this is less of a concern because a significant difference between groups was still detected

P-value: the probability that the observed difference between two groups for a *specific outcome* is due to chance only.

Level of significance (aka alpha): the probability the investigator is willing to take that the results occurred due to chance alone.

- Typically, the level of significance is set at 0.05 (5%) which would indicate that the investigators are willing to accept a 5% chance for a false positive result
- If the p-value is less than alpha then the results are considered statistically significant

Confidence interval: range of values in which the true value likely resides.

- A 95% confidence interval means that if a trial is *repeated* several times from the same sample population then it would be expected that 95% of the confidence intervals would contain the true value for said measure (*and 5% would not*)
- For hazards ratios, if the confidence interval for a measurement includes the value of 1.00 then the difference cannot be considered statistically significant due to a hazards ratio of 1.00 indicating no difference between treatment groups
- The wider the confidence interval the less precise an estimate can be made for the true value of said measure

Non-inferiority trial: designed to assess if the active treatment is no worse than the control treatment by a predetermined margin (aka non-inferiority margin).

- A non-inferiority trial cannot be used to claim superiority without predetermined testing specified in the protocol
- Please note - superiority trials cannot be used to claim non-inferiority

Non-inferiority margin (aka NI margin): the difference allowed between active and control treatments to be considered non-inferior.

- If the NI margin is set at 1.30 then to meet criteria for non-inferiority the confidence interval for said measure must not include 1.30
- If the confidence interval crosses/touches the NI margin then non-inferiority cannot be claimed

Intent to treat population (ITT): the sample of patients that underwent randomization into the trial.

Modified intent to treat population (mITT): the sample of patients that underwent randomization and met one or more qualifying criteria.

- Typically, the additional criteria is receiving at least one dose of study medication in order to be included in a safety analysis

Per protocol population (PP): the sample of patients that successfully completed the trial.

Composite endpoints: a combination of outcomes reported for a single measure of effect.

- Example: composite of cardiovascular death, myocardial infarction or stroke
- Each component should ideally occur at similar rates and have similar clinical significance to avoid distortion of the overall composite measure
- A composite endpoint of cardiovascular death or minor bleeding would not be appropriate due to death being much more significant than minor bleeding

Relative risk reduction: the change in event rate of the active group relative to the control group.

- $RRR = 1 - (\text{active/control})$
- Relative risk reduction is more commonly used when reporting treatment effect
 - However, it is subject to misinterpretation and overestimation of treatment effect
- For example, if the event rate in group A was 10% and group B was 20% this would represent a 50% relative risk reduction but only a true treatment difference of 10%

Absolute risk reduction: the absolute change in event rates between two groups.

- $ARR = \text{control event rate} - \text{active event rate}$
- Less commonly reported when reporting treatment effect

Number needed to treat: an estimate of how many patients would need to receive “Treatment A” to prevent one outcome compared to “Treatment B”.

- $NNT = 1/ARR$ (absolute risk reduction)
- Please note - that NNT must be reported as a whole integer (rounded up)
- It is important to consider the time frame of the trial when interpreting the NNT
- For example, a trial averaging 3 years with a NNT of 14 should be interpreted as that for *approximately* every 14 patients given “Therapy A” for an *average* of 3 years one clinical outcome would be prevented compared to patients receiving “Therapy B”

Number needed to harm: an estimate of how many patients would need to receive “Treatment A” for one adverse outcome to occur compared to “Treatment B”.

- $NNH = 1/ARR$ (absolute risk reduction)
- Please note - that NNH must be reported as a whole integer (rounded down)
- It is important to consider the time frame of the trial when interpreting the NNH
- For example, a trial averaging 3 years with a NNH of 7 should be interpreted as that for *approximately* every 7 patients given “Treatment A” for an *average* of 3 years one adverse clinical outcome would occur compared to patients receiving “Therapy B”

While NNT and NNH are simple to calculate and appear straightforward to use, it is very important to remember that these values are *estimates* based on trial results used to help illustrate the magnitude of treatment effect in terms of patients instead of percentages.

- A NNT value lower than a NNH value indicates that the benefit/risk ratio is favorable, however these calculations are based off average trial results from populations that may differ significantly from a specific patient
- Additionally, the clinical significance of each outcome must be considered (example - cardiovascular death vs hypotension)
- It is only appropriate to report NNT/NNH for statistically significant differences

Level of evidence: the measure of the quality of evidence from a trial.

- Level I - randomized, controlled trial with power set and met
- Level II - randomized, controlled trial with power set but not met
- Levels III, IV or V - observational trials with or without a control group

Grade of recommendation: used to rate the strength of a recommendation.

- The higher the level of evidence the higher the grade of recommendation
 - Level I - Grade A
 - Level II - Grade B
- However, depending on other factors and considerations a higher or lower recommendation may be given for a trial

Additional Information

The following resources were used to form this quick biostatistics review:

- Malone PM, Witt BA, Malone MJ, Peterson DM. eds. *Drug Information: A Guide for Pharmacists, Seventh edition*. McGraw-Hill Education; 2022.
- Bryant PJ, Pace HA. *The Pharmacist's Guide to Evidence-Based Medicine for Clinical Decision Making*. American Society of Health-System Pharmacists; 2008.

Please refer to these resources for more thorough and comprehensive information on the subject.

4S

Randomized trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet*. 1994;344(8934):1383-1389.

Objective: To determine the effect of simvastatin on morbidity and mortality outcomes in patients with coronary heart disease and elevated cholesterol levels.

Primary Efficacy Measure: All-cause mortality

Secondary Efficacy Measure: Composite of coronary death, non-fatal myocardial infarction, resuscitated cardiac arrest and silent myocardial infarction

Participants: Patients with history of coronary heart disease

- Age ~59 years; ~81% male
- Qualifying event myocardial infarction only ~62%; angina only ~21%; both ~16%
- Total cholesterol ~260 mg/dL; LDL ~188 mg/dL; HDL ~70 mg/dL
- Triglycerides ~133 mg/dL
- Beta-blocker ~57%; aspirin ~37%; nitrate ~32%; CCB ~31%

Inclusion Criteria:

- Age 35-70 years
- History of acute myocardial infarction or angina pectoris
- Total cholesterol 5.5 to 8.0 mmol/L (~212 mg/dL to ~309 mg/dL)
- Triglycerides \leq 2.5 mmol/L (~221 mg/dL)

Exclusion Criteria:

- Unstable angina
- Myocardial infarction within previous 6 months
- Planned coronary artery surgery or angioplasty
- Antiarrhythmic therapy or persistent atrial fibrillation
- Congestive heart failure requiring digitalis, diuretics or vasodilators
- History of completed stroke

Drug: Simvastatin

Design: Randomized, double-blind, placebo-controlled trial

Methods: Eligible patients with serum total cholesterol $>$ 5.5 mmol/L (~212 mg/dL) were given dietary advice and after 8 weeks a second blood draw was performed and a 2-week placebo run-in period was initiated. Patients meeting the cholesterol and triglyceride requirements were randomized to receive either simvastatin 20 mg once daily before evening meal or matching placebo. Dosages could be adjusted at the 12 week and 6 month visits based on prior serum cholesterol values. The goal of simvastatin therapy was to reduce total cholesterol to 3.0-5.2 mmol/L (~116 to ~201 mg/dL). If patients were out of range then the dosage was increased to 40 mg daily or reduced to 10 mg daily.

Duration: Median follow-up period of 5.4 years

Statistical Analysis: It was determined that 4400 randomized patients and 440 deaths would achieve 95% power ($\alpha = 0.05$). The ITT population was used for all analyses.

Results: A total of 4444 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. The trial was stopped early based on recommendation from the safety and monitoring committee due to interim analysis demonstrating that the pre-specified boundary had been crossed in favor of simvastatin. In the simvastatin group, 37% of patients had their dosage raised to 40 mg daily during the first six months (only 2 patients were reduced to 10 mg daily). After one year, 72% of simvastatin patients were within the target range for total cholesterol. Coronary death accounted for 74% of the total mortality events in this trial.

Placebo (N=2223) Vs Simvastatin (N=2221)

All-Cause Mortality:

256 (11.5%) vs 182 (8.19%); HR 0.70 (95% CI 0.58-0.85)
p=0.0003; ARR 3.32%; NNT ~31

Coronary Death:

189 (8.50%) vs 111 (5.00%); HR 0.58 (95% CI 0.46-0.73)

Secondary Composite Outcome:

622 (28.0%) vs 431 (19.4%); HR 0.66 (95% CI 0.59-0.75)
p<0.00001; ARR 8.57%; NNT ~12

Limitations:

- External validity - trial conducted entirely in Scandinavia
- Patient population - cannot extrapolate results to patients without history of coronary artery disease

Level of Evidence: Level I - with minor limitations

Recommendation: For these reasons, I recommend the use of simvastatin 20-40 mg once daily to reduce cardiovascular morbidity and mortality in patients with coronary heart disease and elevated cholesterol levels.

Efficacy:

- All-cause mortality was significantly lower in the simvastatin group compared to placebo (driven primarily by reduced rates of coronary death)
- The rate of the secondary composite outcome was also significantly lower in the simvastatin group

Safety:

- There was one case of rhabdomyolysis in simvastatin group (full recovery)
- Rates of AST liver enzyme elevation (3x ULN) were similar between treatment groups
- ALT liver enzyme elevation (3x ULN) occurred more often in the simvastatin group compared to placebo (49 patients vs 33 patients, respectively)

Cost:

- The cost of using simvastatin must be balanced against the cost-savings of reducing cardiovascular morbidity and mortality

Special Considerations/Populations:

- Patient population must be considered - all had history of coronary heart disease and elevated cholesterol levels

Grade of Recommendation: A

ACCORD-BG

Jamerson K, Weber MA, Bakris GL, et al. Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high-risk patients. *N Engl J Med.* 2008;359(23):2417-2428.

Objective: To determine the effect of intensive glycemic control compared to standard glycemic control on cardiovascular event rates in patients with type 2 diabetes and cardiovascular disease or risk-factors.

Primary Efficacy Measure: Composite of cardiovascular death, non-fatal myocardial infarction and non-fatal stroke

Secondary Efficacy Measure: All-cause mortality

Participants: Patients with type 2 diabetes and established cardiovascular disease or risk-factors

- Age ~62 years; male ~62%
- HgA1c ~8.3%; fasting blood glucose ~175 mg/dL; previous cardiovascular event ~35%
- Statin ~62%; metformin ~60%; sulfonylurea ~50%; aspirin ~54%

Inclusion Criteria:

- Patients with type 2 diabetes and HgA1c $\geq 7.5\%$ plus one of the following:
 - Age 40-79 with established cardiovascular disease
 - Age 55-79 with anatomical evidence of atherosclerosis, albuminuria, LV hypertrophy, OR ≥ 2 additional risk factors for cardiovascular disease (dyslipidemia, hypertension, current smoker or obesity)

Exclusion Criteria:

- Frequent or recent serious hypoglycemic events
- Unwilling to perform home blood glucose monitoring
- BMI > 45
- SCr > 1.5 mg/dL

Drugs: n/a

Design: Randomized, open-label, active-comparison trial

Methods: Eligible patients were randomized to either intensive therapy (target HgA1c < 6%) or standard therapy (target HgA1c 7.0-7.9%). Patients were provided glucose-lowering medications from the trial formulary, however the use of outside medications to treat hyperglycemia was allowed. Patient medication regimens were determined individually based on treatment group and response to therapy.

Duration: Mean follow-up period of 3.5 years

Statistical Analysis: It was determined that a follow-up period of at least 5.6 years and an event rate $\geq 2.9\%$ for the primary composite outcome would achieve 89% power ($\alpha = 0.05$). The ITT population was used for the primary and secondary efficacy analyses.

Results: A total of 10,521 patients underwent randomization. The trial was stopped early at the recommendation of the safety committee due to data showing increased mortality rates in the intensive therapy group. After one year of follow-up, the median HgA1c for the intensive and standard treatment groups was 6.4% and 7.5%, respectively. These HgA1c levels were maintained throughout the follow-up period. The intensive treatment group had greater exposure to medications (from all classes) as well as significantly higher rates of adverse events (hypoglycemia, weight gain and fluid retention). The occurrence of weight gain greater than 10 kg from baseline was 27.8% in the intensive therapy group and 14.1% in the standard therapy group ($p < 0.001$).

Intensive Treatment (N=5128) Vs Standard Treatment (N=5123)

Composite of Cardiovascular Death, Non-Fatal Myocardial Infarction & Non-Fatal Stroke:

352 (6.86%) vs 371 (7.24%); HR 0.90 (95% CI 0.78-1.04); p=0.16

All-Cause Mortality:

257 (5.01%) vs 203 (3.96%); HR 1.22 (95% CI 1.01-1.46)
p=0.04; ARI 1.05%; NNT ~96

Cardiovascular Death:

135 (2.63%) vs 94 (1.83%); HR 1.35 (95% CI 1.04-1.76)
p=0.02; ARI 0.80%; NNH ~126

Non-Fatal Myocardial Infarction:

186 (3.63%) vs 235 (4.59%); HR 0.76 (95% CI 0.62-0.92)
p=0.004; ARR 0.96%; NNT ~105

Non-Fatal Stroke:

67 (1.31%) vs 61 (1.19%); HR 1.06 (95% CI 0.75-1.50); p=0.74

Hypoglycemia Requiring Medical Assistance:

538 (10.5%) vs 179 (3.49%)
p<0.001; ARI 7.01%; NNH ~14

Limitations:

- Power set but not met due to the trial being stopped early (clinical significance likely low)
 - It is possible that the trial duration (~3.5 years) was not sufficient to demonstrate the full cardiovascular benefit of intensive therapy (although it was sufficient to demonstrate several safety concerns)
- Open-label trial design

Level of Evidence: Level II - with major limitations

Recommendation: For these reasons, I do not recommend targeting an HgA1c < 6% over standard blood glucose control to reduce the risk of cardiovascular events in high-risk patients with type 2 diabetes.

Efficacy:

- There was no significant difference between treatment groups regarding the primary composite outcome
- Rates of cardiovascular death and all-cause mortality were significantly higher in the intensive treatment group

Safety:

- Rates of hypoglycemia requiring medical assistance were significantly higher in the intensive therapy group
- Fluid retention and weight gain (> 10 kg) occurred at significantly higher rates in the intensive therapy group

Cost:

- Any potential benefit of intensive glycemc therapy must be balanced against the increased cost of achieving the lower target HgA1c (via increased medication usage) as well as the added costs of monitoring and managing hypoglycemic episodes

Special Considerations/Populations:

- This trial demonstrated greater harm than benefit with intensive therapy
- Results cannot be applied to a younger, less complicated patient population

Grade of Recommendation: B

ALLHAT

ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA*. 2002;288(23):2981-2997.

Objective: To determine the effect of CCBs, ACEis and alpha-blockers compared to thiazide-type diuretics on morbidity and mortality in high-risk patients with hypertension.

Primary Efficacy Measure: Composite of fatal coronary heart disease or non-fatal myocardial infarction

Secondary Efficacy Measures: All-cause mortality, fatal/non-fatal stroke, combined coronary heart disease, combined cardiovascular disease

Participants: Patients with stage 1 or 2 hypertension at high-risk for cardiac events

- Age ~67 years; male ~53%
- BP ~146/84 mmHg
- Baseline ASCVD ~52%

Inclusion Criteria:

- Age \geq 55 years
- Hypertension (stage 1 or 2) plus \geq 1 risk factor for coronary heart disease (previous myocardial infarction or stroke $>$ 6 months prior, left-ventricular hypertrophy, type 2 diabetes, current cigarette smoker, HDL $<$ 35 mg/dL, documented ASCVD)

Exclusion Criteria:

- History of hospitalization due to heart failure
- Symptomatic heart failure
- LVEF $<$ 35%

Drugs: Chlorthalidone; lisinopril, amlodipine, doxazosin

Design: Randomized, double-blind, active-comparison trial

Methods: Eligible patients were allowed to continue previous medications until they received the study drug, at which point all previous medications were stopped. Study medication was titrated to a target blood pressure of $<$ 140/90 mmHg in all groups (chlorthalidone 12.5 – 25 mg; amlodipine 2.5 – 10 mg; lisinopril 10 – 40 mg). The use of open-label agents was allowed per investigator discretion.

Amlodipine represented the CCB arm, lisinopril the ACEi arm and chlorthalidone the diuretic arm. The doxazosin treatment arm was stopped early due to increased rates of heart failure (not included in this analysis). Patients underwent randomization at a ratio of 1.7:1:1 (higher for the diuretic arm to maximize power).

Duration: Mean follow-up period of 4.9 years

Statistical Analysis: It was determined that 40,000 randomized patients would provide ~83% for the primary analysis ($\alpha = 0.0178$). For analysis of secondary outcomes, pre-specified components of combined outcomes and patient subgroups a p-value $<$ 0.05 was considered statistically significant. The ITT population was used for analyses.

Results: A total of 42,418 patients underwent randomization (9061 patients randomized to doxazosin group). No significant difference was seen for the primary outcome for either amlodipine or lisinopril compared to chlorthalidone.

There was no significant difference for any secondary outcomes between amlodipine and chlorthalidone, however the pre-specified individual components of the combined secondary outcomes were examined and significant differences were seen for heart failure and heart failure hospitalizations/deaths favoring chlorthalidone. Average SBP at 5 years was significantly lower in the chlorthalidone group compared to amlodipine (133.9 mmHg vs 134.7 mmHg; p=0.03).

Amlodipine (N=9048) Vs Chlorthalidone (N=15,255)

Composite of Fatal Coronary Heart Disease & Non-Fatal Myocardial Infarction:

798 (8.82%) vs 1362 (8.93%); RR 0.98 (95% CI 0.90-1.07); p=0.65

Non-fatal myocardial infarction was ~64-66% of composite outcome

Heart Failure:

706 (7.80%) vs 870 (5.70%); RR 1.38 (95% CI 1.25-1.52)

p<0.001; ARR 2.10%; NNT ~48

Heart Failure Hospitalizations/Heart Failure Death:

578 (6.39%) vs 724 (4.75%); RR 1.35 (95% CI 1.21-1.50)

p<0.001; ARR 1.64%; NNT ~61

There were significant differences for the secondary outcomes of stroke and combined cardiovascular disease between lisinopril and chlorthalidone favoring the thiazide group. For the pre-specified individual components of the combined secondary outcomes, significant differences were seen in heart failure and angina hospitalizations/treatment favoring chlorthalidone. Average SBP at 5 years was significantly lower in the chlorthalidone group (133.9 mmHg vs 135.9 mmHg; p<0.001).

Lisinopril (N=9054) Vs Chlorthalidone (15,255)

Composite of Fatal Coronary Heart Disease & Non-Fatal Myocardial Infarction:

796 (8.79%) vs 1362 (8.93%); RR 0.99 (95% CI 0.91-1.08); p=0.81

Non-fatal myocardial infarction was ~64-66% of composite outcome

Stroke:

457 (5.05%) vs 675 (4.42%); RR 1.15 (95% CI 1.02-1.30)

p=0.02; ARR 0.62%; NNT ~161

Combined Cardiovascular Disease:

2514 (27.8%) vs 3941 (25.8%); RR 1.10 (95% CI 1.05-1.16)

p<0.001; ARR 1.93%; NNT ~52

Heart Failure:

612 (6.76%) vs 870 (5.70%); RR 1.19 (95% CI 1.07-1.31)

p<0.001; ARR 1.06%; NNT ~95

Hospitalized/Treated Angina:

1019 (11.3%) vs 1567 (10.3%); RR 1.11 (95% CI 1.03-1.20)

p=0.01; ARR 0.98%; NNT ~102

Limitations:

- Use of open-label add-on antihypertensive therapy is a potential confounding factor (however, statistical differences still seen between treatment groups for multiple outcomes)

Level of Evidence: Level I - with minor limitations

Recommendation: For these reasons, I recommend the use of chlorthalidone over amlodipine and lisinopril as the preferred first-line therapy for high-risk hypertension patients. However, the selection of initial antihypertensive therapy must be individualized according to patient-specific characteristics.

Efficacy:

- There was no significant difference in the rates of the primary composite outcome of fatal coronary heart disease and non-fatal myocardial infarction between chlorthalidone, amlodipine and lisinopril
- Rates of heart failure and heart failure hospitalizations/death were significantly lower in the chlorthalidone group compared to the amlodipine group
- Rates of stroke, combined cardiovascular disease, heart failure and hospitalized/treated angina were significantly lower in the chlorthalidone group compared to the lisinopril group
- Average SBP at 5 years was significantly lower in the chlorthalidone group compared to the amlodipine and lisinopril groups

Safety:

- There were significantly higher rates of angioedema in the lisinopril group compared to the chlorthalidone group
- Chlorthalidone demonstrated significant biochemical changes (increased cholesterol and fasting glucose levels, decreased potassium levels) compared to lisinopril and amlodipine
 - However, these are known and predictable effects of this medication class that can be monitored
 - Additionally, these biochemical changes did not yield net increases in negative cardiovascular outcomes compared to the other treatment arms

Cost:

- The cost of using chlorthalidone over lisinopril or amlodipine must be balanced against the cost-savings of preventing cardiovascular outcomes (specifically, heart failure events)

Special Considerations/Populations:

- Cannot extrapolate the treatment effect of chlorthalidone to other diuretics such as hydrochlorothiazide (different classes of thiazide with different properties)

Grade of Recommendation: A

DAPA-HF

McMurray JJV, Solomon SD, Inzucchi SE, et al. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med.* 2019;381(21):1995-2008.

Objective: To determine the effect of dapagliflozin on morbidity and mortality outcomes in heart failure patients with reduced ejection fraction with or without type 2 diabetes.

Primary Efficacy Measure: Composite of worsening heart failure (heart failure hospitalization or urgent visit requiring IV treatment for heart failure) and cardiovascular death

Participants: Patients with heart failure and reduced ejection fraction (with or without type 2 diabetes)

- Age ~66 years; male ~77%;
- NYHA class II ~67%; class III ~31%; class IV ~1%
- LVEF ~31%; HR ~72; eGFR ~66 mL/min
- History of type 2 diabetes ~42%
- Baseline ACEi ~56%; ARB ~27%; ARNi ~11%; beta-blocker ~96%; MRA ~71%

Inclusion Criteria:

- Age \geq 18 years
- LVEF \leq 40%
- NYHA class II-IV
- NT-proBNP \geq 600 pg/mL (\geq 400 pg/mL if hospitalized for heart failure within prior year)
- Receiving standard heart failure therapy (ACEi/ARB/ARNi plus beta-blocker)

Exclusion Criteria:

- Type 1 diabetes
- eGFR $<$ 30 mL/min
- Symptoms of hypotension or SBP $<$ 95 mmHg

Drug: Dapagliflozin

Design: Randomized, double-blind, placebo-controlled trial

Methods: Eligible patients were randomized to receive either dapagliflozin 10 mg daily or matching placebo. Standard heart failure (and type 2 diabetes) treatment was continued and adjusted if needed.

Duration: Median follow-up period of 18.2 months

Statistical Analysis: It was determined that 844 primary events were required to achieve 90% power ($\alpha = 0.05$). The ITT population was used for all efficacy analyses.

Results: A total of 4744 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. The average HgA1c decreased by a significantly greater amount in the dapagliflozin group compared to placebo (-0.21% vs +0.04%; $p < 0.001$). The treatment effect of dapagliflozin was consistent in patients with and without type 2 diabetes.

Dapagliflozin (N=2373) Vs Placebo (N=2371)

Primary Composite Outcome:

386 (16.3%) vs 502 (21.2%); HR 0.74 (95% CI 0.65-0.85)
p<0.001; ARR 4.91%; NNT ~21

Heart Failure Hospitalization:

231 (9.73%) vs 318 (13.4%); HR 0.70 (95% CI 0.59-0.83)

Urgent Visit Requiring IV Treatment for Heart Failure:

10 (0.42%) vs 23 (0.97%); HR 0.43 (95% CI 0.20-0.90)

Cardiovascular Death:

227 (9.57%) vs 273 (11.5%); HR 0.82 (95% CI 0.69-0.98)

Limitations:

- External validity - cannot extrapolate results to patients with preserved ejection fraction

Level of Evidence: Level I - with minor limitations

Recommendation: For these reasons, I recommend the use of dapagliflozin 10 mg to reduce morbidity and mortality rates in heart failure reduced ejection fraction patients with or without type 2 diabetes. However, it would be reasonable to optimize standard heart failure therapy prior to initiation of dapagliflozin to maximize overall treatment benefit.

Efficacy:

- The dapagliflozin treatment group demonstrated significantly lower rates of the primary composite outcome compared to placebo (both individual components favored dapagliflozin significantly)
- The treatment benefit of dapagliflozin was consistent in patients with and without type 2 diabetes

Safety:

- There were no significant differences in rates of volume depletion, renal adverse events, fractures, amputations, major hypoglycemia, or gangrene
- All and major hypoglycemia events occurred in patients with type 2 diabetes

Cost:

- The cost of using dapagliflozin must be balanced against the cost-savings of preventing worsening heart failure and cardiovascular death

Special Considerations/Populations:

- Results must be considered as dapagliflozin in addition to standard heart failure therapy
- Cannot extrapolate data to patients with heart failure and preserved ejection fraction

Grade of Recommendation: A

PARADIGM-HF

McMurray JJ, Packer M, Desai AS, et al. Angiotensin-nepriylsin inhibition versus enalapril in heart failure. *N Engl J Med.* 2014;371(11):993-1004.

Objective: To determine the effect of sacubitril-valsartan compared to enalapril on morbidity and mortality outcomes in patients with heart failure and reduced ejection fraction.

Primary Efficacy Measure: Composite of cardiovascular death or first heart failure hospitalization

Participants: Patients with heart failure and reduced ejection fraction (NYHA class II-IV)

- Age ~64 years; male ~79%
- LVEF ~30%
- NYHA class II ~70%; class III ~24%
- SBP ~122 mmHg; HR ~72 bpm
- Baseline beta-blocker ~93%; diuretic ~80%; mineralocorticoid antagonist ~55%

Inclusion Criteria:

- Age ≥ 18 years with LVEF $\leq 35\%$ (originally 40%)
- NYHA functional class II-IV
- BNP ≥ 150 pg/mL (≥ 100 pg/mL if hospitalized for heart failure within previous 12 months)
- Receiving stable dosing of beta-blocker and ACEi/ARB for 4 weeks

Exclusion Criteria:

- Symptomatic hypotension
- SBP < 100 mmHg
- eGFR < 30 mL/min
- Serum potassium > 5.2 mmol/L
- History of angioedema

Drugs: Sacubitril/valsartan; enalapril

Design: Randomized, double-blind, active-controlled trial

Methods: Eligible patients underwent two single-blind run-in phases. First, all patients received enalapril 10 mg twice daily alone for 2 weeks and then all patients received ARNi therapy (sacubitril/valsartan) for 4-6 weeks (initially 49 mg/51 mg twice daily, then increased to 97 mg/103 mg twice daily) alone to ensure the side effect profile was acceptable. Those that successfully completed these run-in periods were then randomized to receive enalapril 10 mg twice daily or sacubitril-valsartan 97 mg/103 mg twice daily. Dosing could be reduced if appropriate.

Duration: Median follow-up period of 27 months

Statistical Analysis: It was determined that 2410 primary events would achieve 97% power ($\alpha = 0.05$). Criteria for stopping the trial early was established by the safety and monitoring committee. The ITT population was used for the efficacy analyses.

Results: A total of 8399 patients underwent randomization. Baseline characteristics were similar between treatment groups. The trial was stopped early due to the results of the third interim analysis showing clear benefit of sacubitril/valsartan over enalapril. The average daily dosing of the study drugs was ~375 mg sacubitril/valsartan and ~19 mg enalapril. At 8 months, the decline in average KCCQ clinical summary score (used to subjectively assess heart failure symptoms and physical limitations) was significantly less in the sacubitril/valsartan group compared to enalapril (-2.99 vs -4.63; $p=0.001$). There was no significant difference in rates of angioedema.

Sacubitril/Valsartan (N=4187) Vs Enalapril (N=4212)

Composite of Cardiovascular Death or First Heart Failure Hospitalization:

914 (21.8%) vs 1117 (26.5%); HR 0.80 (95% CI 0.73-0.87)
p<0.001; ARR 4.69%; NNT ~22

Cardiovascular Death:

558 (13.3%) vs 693 (16.5%); HR 0.80 (95% CI 0.71-0.89)
p<0.001; ARR 3.13%; NNT ~32

First Heart Failure Hospitalization:

537 (12.8%) vs 658 (15.6%); HR 0.79 (95% CI 0.71-0.89)
p<0.001; ARR 2.80%; NNT ~36

All-Cause Mortality:

711 (17.0%) vs 835 (19.8%); HR 0.84 (95% CI 0.76-0.93)
p<0.001; ARR 2.84%; NNT ~36

Symptomatic Hypotension with SBP < 90 mmHg:

112 (2.67%) vs 59 (1.40%); p<0.001; ARI ~1.27%; NNH ~78

Limitations:

- Power set but not met - failed to achieve 2410 primary event outcomes (trial stopped early - clinical significance minimal)
- Patient population - cannot extrapolate trial results to those with preserved ejection fraction

Level of Evidence: Level II - with major limitations

Recommendation: For these reasons, I recommend the use of sacubitril/valsartan over enalapril to further reduce morbidity and mortality rates in heart failure patients with reduced ejection fraction. However, it would be reasonable to optimize standard heart failure therapy prior to initiation of sacubitril/valsartan to maximize the overall treatment benefit.

Efficacy:

- Rates of the primary composite outcome were significantly lower in the sacubitril/valsartan group compared to enalapril
- The individual components of cardiovascular death and first heart failure hospitalization were significantly lower in the sacubitril/valsartan group as well
- Average KCCQ score decline was significantly less in the sacubitril/valsartan group compared to enalapril (indicates less decline in quality of life)

Safety:

- There was no significant difference in the rates of angioedema
- Symptomatic hypotension (with SBP < 90 mmHg) did occur at significantly higher rates in the sacubitril/valsartan group compared to enalapril

Cost:

- The cost of using sacubitril/valsartan must be balanced against the cost-savings of preventing morbidity and mortality events in heart failure patients

Special Considerations/Populations:

- If switching from ACEi to ARNi (or vice versa) there must be a wash-out period of 36 hours to reduce the risk for angioedema
- It is important to note that all patients included in this trial were able to tolerate ACEi prior to starting ARNi therapy (due to trial design)

Grade of Recommendation: A

PRECISION

Nissen SE, Yeomans ND, Solomon DH, et al. Cardiovascular Safety of Celecoxib, Naproxen, or Ibuprofen for Arthritis. *N Engl J Med.* 2016;375(26):2519-2529.

Objective: To determine the effect of celecoxib on cardiovascular outcomes compared to ibuprofen and naproxen in patients with arthritis pain.

Primary Safety Measure: Composite of cardiovascular death, non-fatal myocardial infarction or non-fatal stroke

Participants: Patients with arthritic pain at increased risk for cardiovascular event

- Age ~63 years; male ~35%
- Osteoarthritis ~90%
- Established cardiovascular disease ~23%; increased risk for cardiovascular event ~77%

Inclusion Criteria:

- Age \geq 18 years
- Required daily treatment with NSAIDs for arthritis pain
- Established cardiovascular disease or increased risk for cardiovascular disease

Exclusion Criteria:

- Arthritis pain controlled adequately by acetaminophen

Drugs: Celecoxib; ibuprofen; naproxen

Design: Randomized, double-blind, active-controlled, non-inferiority trial

Methods: Eligible patients were randomized 1:1:1 to receive celecoxib 100 mg twice daily, ibuprofen 600 mg three times daily or naproxen 375 mg twice daily (plus matching placebo). Doses could be increased at the discretion of the investigators to celecoxib 200 mg twice daily, ibuprofen 800 mg three times daily or naproxen 500 mg twice daily. Esomeprazole 20-40 mg daily was used for gastric protection in all patients.

Duration: Mean follow-up period of ~34 months

Statistical Analysis: Naproxen was used as the primary comparator for the non-inferiority analysis. To demonstrate non-inferiority the following criteria were required: HR \leq 1.12 with upper limit of CI $<$ 1.33 in both ITT and on-treatment analysis populations. It was originally determined that 762 primary events would provide 90% for the non-inferiority analysis. However, due to lower than expected event rates it was recommended to amend the protocol to make the upper CI limit $<$ 1.40 and require 580 primary events in the ITT population and 420 events in the on-treatment population to provide 80% power with a minimum of 18 months of follow-up. A p-value $<$ 0.05 was considered statistically significant for the secondary safety measure and a p $<$ 0.025 for the primary safety measure.

Results: A total of 24,222 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. The mean daily doses of the study medications were ~209 mg celecoxib, ~852 mg naproxen and ~2045 mg ibuprofen. Celecoxib demonstrated non-inferiority to naproxen in both the ITT and on-treatment analysis. Visual analog scale analysis showed statistically significant benefit of naproxen over both celecoxib and ibuprofen for pain control (p $<$ 0.001 and 0.01, respectively). However, the difference in the VAS was not considered clinically significant due to being $<$ 13.7 mm. However, There was no significant difference in VAS measurements between celecoxib and ibuprofen (p=0.38).

Celecoxib (N=8072) Vs Naproxen (N=7969)

Cardiovascular Death, Non-Fatal Myocardial Infarction or Non-Fatal Stroke:

188 (2.33%) vs 201 (2.52%); HR 0.93 (95% CI 0.76-1.13); p=0.45

Rates of the individual components were not significantly different

Serious Gastrointestinal Events:

86 (1.07%) vs 119 (1.49%); HR 0.71 (95% CI 0.54-0.93)

p=0.01; ARI 0.43%; NNH ~223

Celecoxib (N=8072) Vs Ibuprofen (N=8040)

Cardiovascular Death, Non-Fatal Myocardial Infarction or Non-Fatal Stroke:

188 (2.33%) vs 218 (2.71%); HR 0.85 (95% CI 0.70-1.04); p=0.12

Rates of the individual components were not significantly different

Serious Gastrointestinal Events:

86 (1.07%) vs 130 (1.62%); HR 0.65 (95% CI 0.50-0.85)

p=0.002; ARI 0.57%; NNH ~176

Limitations:

- Use of esomeprazole in all patients likely reduced the overall rates of serious gastrointestinal events, however its use is reasonable to limit patient risk
- Patient population must be considered - vast majority of patients did not have established cardiovascular disease at baseline

Level of Evidence: Level I - with minor limitations

Recommendation: For these reasons, I recommend the use of celecoxib as an alternative to non-selective NSAIDs in arthritis patients at increased risk for cardiovascular events, especially in those with increased risk for gastrointestinal bleeds.

Efficacy:

- Celecoxib demonstrated non-inferiority to ibuprofen and naproxen regarding the the cardiovascular composite outcome
- There was no significant difference in the rates of the primary composite outcome or the individual components between treatment groups

Safety:

- Rates of serious gastrointestinal events were significantly lower in the celecoxib group compared to naproxen and ibuprofen groups (despite use of esomeprazole in all patients)

Cost:

- The cost of using celecoxib must be balanced against the cost of using naproxen or ibuprofen
 - However, the cost-savings of avoiding serious gastrointestinal events must be considered

Special Considerations/Populations:

- Celecoxib is a COX-2 selective inhibitor and thus less likely to impact gastric mucosa compared to non-selective NSAIDs
- The use of daily esomeprazole must be considered when interpreting safety data

Grade of Recommendation: A

RACE II

Van Gelder IC, Groenveld HF, Crijns HJ, et al. Lenient versus strict rate control in patients with atrial fibrillation. *N Engl J Med.* 2010;362(15):1363-1373.

Objective: To determine the effect of lenient rate control compared to strict rate control on clinical outcomes in patients with atrial fibrillation.

Primary Efficacy Measure: Composite of cardiovascular death, heart failure hospitalization, stroke, systemic embolism, major bleeding, arrhythmic events, sustained ventricular tachycardia, cardiac arrest, life-threatening adverse drug reactions due to rate control medications or pacemaker implantation

Participants: Patients with atrial fibrillation

- Age ~68 years; male ~66%
- CHADS₂ score 0-1 ~61%; score = 2 ~26%; score 3-6 ~13%
- BP ~136/93 mmHg; HR ~96 bpm
- Managed on beta-blocker alone ~45%

Inclusion Criteria:

- Permanent atrial fibrillation for up to 12 months
- Age ≤ 80 years
- Average resting HR > 80 bpm
- Current use of oral anticoagulant therapy (or aspirin)

Exclusion Criteria:

- Paroxysmal atrial fibrillation
- Contraindications for strict or lenient rate control
- Unstable heart failure
- History of any stroke
- Cardiac surgery within previous 3 months

Treatment: Lenient rate control (resting HR < 110 bpm); strict rate control (resting HR < 80 bpm)

Design: Randomized, open-label, active-controlled, non-inferiority trial

Methods: Eligible patients were randomized to receive strict rate control (target resting HR < 80 bpm) or lenient rate control (target resting HR < 110 bpm). Beta-blockers, non-DHP calcium channel blockers and digoxin were to be used alone or in combination to achieve the target heart rate.

Duration: Follow-up period of 2 to 3 years

Statistical Analysis: It was determined that 250 randomized patients per treatment group would provide 80% power ($\alpha = 0.05$). To demonstrate non-inferiority the upper limit of the 90% confidence interval for the absolute difference between treatment groups needed to be < 10%. However, due to changes in the treatment period the estimated cumulative incidences at 3 years was used to assess for non-inferiority. The ITT population was used for the primary analyses.

Results: A total of 614 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. In the lenient rate control group ~98% of patients achieved target heart rate compared to ~67% in the strict rate control group. In the lenient control group the majority of patients (~42%) were managed on a beta-blocker alone whereas the majority of the strict control group (~37%) required a beta-blocker plus digoxin. Additionally, the average beta-blocker dose was significantly lower in the lenient rate control group ($p < 0.001$). Average heart rates at the end of the follow-up period were 85 bpm in the lenient rate control group and 76 bpm in the strict rate control group ($p < 0.001$). There was no significant difference in rates of the individual components of the composite outcome.

Lenient Rate Control (N=311) Vs Strict Rate Control (N=303)

Primary Composite Outcome:

38 (12.2%) vs 43 (14.2%); HR 0.84 (90% CI 0.58-1.21)

*Lenient rate control demonstrated non-inferiority compared to strict rate control
Absolute rate difference -2.0% (90% CI -7.6% to 3.5%)*

Limitations:

- Open-label trial design
- Trial population size relatively small

Level of Evidence: Level I - with major limitations

Recommendation: For these reasons, I recommend the use of lenient rate control over strict rate control in patients with atrial fibrillation.

Efficacy:

- Lenient rate control demonstrated non-inferiority to strict rate control regarding the rates of the primary composite outcome
- Average HR 85 bpm in lenient rate control group (~98% achieved target HR < 110 bpm)
- Average HR 76 bpm in strict rate control group (~67% achieved target HR < 80 bpm)
- The strict rate control group required significantly more medications at higher doses compared to the lenient rate control group

Safety:

- Overall rates of adverse drug reactions due to rate control medications were lower in the lenient rate control group

Cost:

- The cost-savings of using fewer medications at lower dosages for lenient rate control must be considered

Special Considerations/Populations:

- It is possible that the difference in average heart rates between treatment groups was not great enough to demonstrate the true treatment effect of each rate control strategy
 - While lenient rate control demonstrated non-inferiority to strict rate control it is important to note that the average heart rate for the lenient rate control group was 85 bpm (which is closer to goal for strict control than lenient control) - this must be considered when interpreting trial results

Grade of Recommendation: A

ROCKET AF

Patel MR, Mahaffey KW, Garg J, et al. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med.* 2011;365(10):883-891.

Objective: To determine the efficacy and safety of rivaroxaban compared to warfarin for prevention of thromboembolic events in patients with atrial fibrillation.

Primary Efficacy Measure: Composite of total stroke and systemic embolism

Primary Safety Measure: Clinically relevant bleeding

Participants: Patients with non-valvular atrial fibrillation at increased risk for thromboembolic event

- Age ~73 years; male ~60%
- Persistent atrial fibrillation ~80%
- CHADS₂ score ~3.5

Inclusion Criteria:

- Age ≥ 18 years with non-valvular atrial fibrillation
- Moderate-high stroke risk:
 - History of stroke/TIA/systemic embolism, or
 - CHADS₂ score ≥ 2

Exclusion Criteria:

- Mitral valve stenosis
- Planned cardioversion
- Active endocarditis
- Prosthetic heart valve
- Increased bleeding risk

Drugs: Rivaroxaban; warfarin

Design: Randomized, double-blind, active-controlled, non-inferiority trial

Methods: Eligible patients were randomized to receive either rivaroxaban 20 mg (15 mg if CrCl 30-49 mL/min) or warfarin (target INR 2.0-3.0) plus matching placebo. The use of aspirin ≤ 100 mg/day was allowed.

Duration: Median follow-up period of 707 days (~2 years)

Statistical Analysis: It was determined that 14,000 randomized patients and 363 primary events would provide 95% power for non-inferiority. A non-inferiority margin of 1.46 was used (alpha = 0.025). If non-inferiority was achieved, then sequential testing for superiority would then be performed using the as-treated population (received one or more doses of study medication and were followed for events regardless of adherence). The per-protocol population (and ITT population) were used for the primary non-inferiority analysis.

Results: A total of 14,264 patients underwent randomization. Baseline patient characteristics were similar between treatment groups except for previous myocardial infarction (higher in the warfarin treatment group). Patients in the warfarin group were within target INR range ~55% of the time. Non-inferiority was demonstrated in both the per-protocol and ITT population analyses. There were conflicting results between the per-protocol and ITT analyses when testing for superiority. The per-protocol analysis demonstrated superiority of rivaroxaban over warfarin, but the ITT analysis failed to demonstrate a significant difference between treatment groups (p=0.12). However, subgroup analysis of the ITT population demonstrated that while on treatment patients in the rivaroxaban group demonstrated significantly lower rates of the primary outcome. The rates of the individual components of the composite outcome were not reported.

Rivaroxaban (N=6958) Vs Warfarin (N=7004)
Per-Protocol Population

Primary Composite Outcome - Non Inferiority:
188 (2.70%) vs 241 (3.44%); HR 0.79 (95% CI 0.66-0.96); p<0.001

Rivaroxaban (N=7061) Vs Warfarin (N=7082)
As-Treated Population

Primary Composite Outcome - Superiority:
189 (2.68%) vs 243 (3.43%); HR 0.79 (0.65-0.95)
p=0.02; ARR 0.75%; NNT ~133

Rivaroxaban (N=7081) Vs Warfarin (N=7090)
ITT Population

Primary Composite Outcome - Superiority:
269 (3.80%) vs 306 (4.32%); HR 0.88 (95% CI 0.75-1.03); p=0.12

Safety:

Rivaroxaban (N=7111) Vs Warfarin (N=7125)

Clinically Relevant Bleeding:
1475 (20.7%) vs 1449 (20.3%); HR 1.03 (95% CI 0.96-1.11); p=0.44

Fatal Bleeding:
27 (0.38%) vs 55 (0.77%); HR 0.50 (95% CI 0.31-0.79)
p=0.003; ARI 0.39%; NNH ~254

Intracranial Hemorrhage:
55 (0.77%) vs 84 (1.18%); HR 0.67 (95% CI 0.47-0.93)
p=0.02; ARI 0.41%; NNH ~246

Major Bleeding:
395 (5.55%) vs 386 (5.42%); HR 1.04 (95% CI 0.90-1.20); p=0.58

Gastrointestinal Bleeding:
224 (3.15%) vs 154 (2.16%); p<0.001; ARI 0.99%; NNH ~101

Limitations:

- Multiple analyses with differing results depending on which patient population is used - interpret cautiously
- Cannot extrapolate results to other DOACs

Level of Evidence: Level I - with minor limitations

Recommendation: For these reasons, I recommend the use of rivaroxaban over warfarin for prevention of thromboembolic events in high-risk patients with non-valvular atrial fibrillation.

Efficacy:

- Rivaroxaban demonstrated non-inferiority to warfarin regarding the primary composite outcome
- Conflicting results regarding superiority of rivaroxaban over warfarin depending on the patient population used for analyses (per-protocol and as-treated analysis demonstrated superiority; ITT analysis failed to demonstrate superiority)
- Subgroup analysis of ITT population showed that while on-treatment, rates of the primary outcome were significantly lower in the rivaroxaban treatment group

Safety:

- Rates of clinically relevant bleeding and major bleeding were not significantly different between treatment groups
- Rates of fatal bleeding and intracranial hemorrhage were significantly lower in the rivaroxaban group compared to warfarin
- Rates of gastrointestinal bleeding were significantly higher in the rivaroxaban group

Cost:

- The cost of using rivaroxaban must be balanced against the cost of using warfarin and monitoring INR
- The cost-savings of preventing thromboembolic events must also be considered

Special Considerations/Populations:

- Cannot apply results to patients with valvular atrial fibrillation

Grade of Recommendation: A

SPRINT

SPRINT Research Group, Wright JT Jr, Williamson JD, et al. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med.* 2015;373(22):2103-2116.

Objective: To determine the effect of intensive blood pressure control compared to standard blood pressure control on cardiovascular outcomes in patients without diabetes.

Primary Efficacy Measure: Composite of cardiovascular death, myocardial infarction, acute coronary syndrome, stroke or heart failure

Participants: Patients with hypertension without diabetes or previous stroke

- Age ~68 years; male ~64%
- BP ~140/78 mmHg
- Established cardiovascular disease at baseline ~20%
- Framingham 10-year cardiovascular risk score ~25%

Inclusion Criteria:

- Age > 50 years
- SBP 130-180 mmHg
- Increased risk of cardiovascular events (one or more of the following):
 - Clinical/subclinical cardiovascular disease other than stroke
 - Chronic kidney disease (eGFR 20 mL/min to less than 60 mL/min)
 - Framingham 10-year cardiovascular risk score > 15%
 - Age > 75 years

Exclusion Criteria:

- Prior stroke
- Diabetes

Drug: n/a

Design: Randomized, open-label, active-controlled trial

Methods: Eligible patients were randomized to either standard blood pressure therapy (target SBP < 140 mmHg) or intensive pressure therapy (target SBP < 120 mmHg). After randomization, hypertension treatments were adjusted accordingly. Investigators were allowed to prescribe any antihypertensive medication(s) but were encouraged to use those with the strongest evidence in reducing cardiovascular events. Thiazide diuretics were encouraged as first line agents (with preference for chlorthalidone). Medications were adjusted to achieve target blood pressure goals (SBP < 120 mmHg for intensive therapy; SBP 135-139 mmHg for standard therapy). Medication dosages were reduced in the standard therapy group if SBP went < 130 mmHg.

Duration: Median follow-up period of 3.26 years

Statistical Analysis: It was determined that 9250 randomized patients would achieve 88.7% power ($\alpha=0.05$). The ITT population was used for all analyses.

Results: A total of 9361 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. The trial was stopped early at the recommendation of the safety committee due to the demonstrated benefit of intensive therapy over standard therapy. The average SBP was 121.5 mmHg in the intensive therapy group and 134.6 mmHg in the standard therapy group. The average number of antihypertensive medications was ~2.8 in the intensive therapy group and ~1.8 in the standard therapy group.

Intensive Therapy (N=4678) Vs Standard Therapy (N=4683)

Primary Composite Outcome:

243 (5.19%) vs 319 (6.81%); HR 0.75 (95% CI 0.64-0.89)
p<0.001; ARR 1.62%; NNT ~62

Cardiovascular Death:

37 (0.79%) vs 65 (1.39%); HR 0.57 (95% CI 0.38-0.85)
p=0.005; ARR 0.60%; NNT ~168

Myocardial Infarction:

97 (2.07%) vs 116 (2.48%); HR 0.83 (95% CI 0.64-1.09); p=0.19

Acute Coronary Syndrome:

40 (0.86%) vs 40 (0.85%); HR 1.00 (95% CI 0.64-1.55); p=0.99

Stroke:

62 (1.33%) vs 70 (1.49%); HR 0.89 (95% CI 0.63-1.25); p=0.50

Heart Failure:

62 (1.33%) vs 100 (2.14%); HR 0.62 (95% CI 0.45-0.84)
p=0.002; ARR 0.81%; NNT ~124

All-Cause Mortality:

155 (3.31%) vs 210 (4.48%); HR 0.73 (95% CI 0.60-0.90)
p=0.003; ARR 1.17%; NNT ~86

Safety:

Hypotension:

110 (2.35%) vs 66 (1.41%); p=0.001; ARI 0.94%; NNH ~106

Syncope:

107 (2.29%) vs 80 (1.71%); p=0.05

Electrolyte Abnormalities:

144 (3.08%) vs 107 (2.28%); p=0.02; ARI 0.79%; NNH ~126

Acute Kidney Injury/Acute Renal Failure:

193 (4.13%) vs 117 (2.50%); p<0.001; ARI 1.63%; NNH ~61

≥ 30% Reduction in GFR to <60 mL (in patients without baseline CKD):
127/3332 (3.81%) vs 37/3345 (1.11%); HR 3.49 (95% CI 2.44-5.10)
p<0.001; ARI 2.71%; NNH ~37

Limitations:

- Open-label trial design
- Patient population must be considered - patients with diabetes were not included

Level of Evidence: Level I - with major limitations

Recommendation: For these reasons, I recommend the use of intensive blood pressure control (SBP < 120 mmHg) over standard blood pressure control (SBP < 140 mmHg) to reduce cardiovascular morbidity and mortality in patients without diabetes or prior stroke. However, the risk for adverse outcomes due to intensive antihypertensive therapy must be considered for each patient.

Efficacy:

- Rates of the primary composite outcome were significantly lower in the intensive blood pressure group compared to standard therapy
- The individual rates of cardiovascular death and heart failure were significantly lower in the intensive therapy group
- All-cause mortality was significantly lower in the intensive therapy group compared to standard therapy

Safety:

- Adverse drug reactions of hypotension, syncope and electrolyte abnormalities occurred significantly more often in the intensive therapy group
- Acute kidney injury/acute renal failure occurred at significantly higher rates in the intensive therapy group
- Patients without baseline chronic kidney disease were significantly more likely to experience significant decreases in GFR in the intensive therapy group compared to standard therapy

Cost:

- The cost of targeting a lower SBP must be balanced against the cost-savings of preventing cardiovascular morbidity and mortality
 - However, the cost of monitoring and managing changes in electrolytes and renal function must also be considered

Special Considerations/Populations:

- Cannot extrapolate results to patients with prior stroke or diabetes
- Thiazide diuretics (specifically chlorthalidone) were encouraged as first-line agents for managing hypertension

Grade of Recommendation: A

VAST-D

Mohamed S, Johnson GR, Chen P, et al. Effect of Antidepressant Switching vs Augmentation on Remission Among Patients With Major Depressive Disorder Unresponsive to Antidepressant Treatment: The VAST-D Randomized Clinical Trial. *JAMA*. 2017;318(2):132-145.

Objective: To compare the efficacy and safety of different treatment strategies (i.e. switching or augmenting) in patients with depression unresponsive to current therapy.

Primary Efficacy Measure: Remission (QIDS-C16 score ≤ 5 at two consecutive follow-up visits during acute treatment phase)

- QIDS-C16 score ranges from 0-27 (higher score, more severe symptoms of depression)

Secondary Efficacy Measures: (1) QIDS-C16 score reduction $\geq 50\%$ from baseline to week 12 (2) CGI Improvement Scale Rating of 1 (very much improved) or 2 (much improved) at any scheduled visit through week 12

Participants: Patients with major depressive disorder unresponsive to current therapy

- Age ~54 years; male ~85%
- Number of previous antidepressant courses ~2 (median)
- QIDS-C16 ~17

Inclusion Criteria:

- Veterans Health Administration (VHA) patients
- Age ≥ 18 years
- Diagnosis of major depressive disorder (MDD)
- Unresponsive to at least one course of antidepressant therapy
 - 6-8 weeks of SSRI, SNRI or mirtazapine

Exclusion Criteria:

- Suicidal ideation requiring inpatient treatment
- Currently treated with bupropion or any antipsychotic
- Dementia
- History of bipolar disorder, schizophrenia, psychosis
- Seizure disorder

Drugs: Aripiprazole, bupropion SR

Design: Randomized, active-controlled trial

Methods: Eligible patients were randomized to one of three treatment groups. Patients would be switched from their current therapy to bupropion SR (switch group) or have current therapy augmented with bupropion SR (augment-bupropion group) or aripiprazole (augment-aripiprazole group). After randomization, both patient and investigator were informed of the assigned treatment group. Bupropion SR dosing was titrated (as tolerated) from 150 mg daily to 300 mg-400 mg daily. Aripiprazole dosing was titrated (as tolerated) from 2 mg daily to 5 mg-15 mg daily. This acute treatment phase lasted 12 weeks.

Duration: 12 weeks

Statistical Analysis: It was determined that 1518 randomized patients would achieve 90% power for the primary efficacy measure comparing the augmented-aripiprazole group to the switch group ($\alpha=0.05$) and the augmented-bupropion group to the switch group ($\alpha=0.025$). A level of significance of 0.025 was used for the secondary efficacy measures.

Results: A total of 1522 patients underwent randomization. Baseline patient characteristics were similar between treatment groups. Both switch and augment-bupropion groups achieved the max dose of bupropion SR 200 mg BID at 12 weeks. The augment-aripiprazole group achieved aripiprazole 10 mg daily at 12 weeks. There was no significant difference in serious adverse event rates between treatment groups. Anxiety occurred at significantly lower rates in the augment-aripiprazole group. However, rates of weight gain ($\geq 7\%$), somnolence and extrapyramidal symptoms were significantly higher in the augment-aripiprazole group compared to the switch group and augment-bupropion group.

Switch Group (N=511) Vs Augment-Bupropion Group (N=506)

Remission:

114 (22.3%) vs 136 (26.9%); RR 1.20 (95% CI 0.97-1.50); p=0.09

$\geq 50\%$ Reduction in QIDS-C16 Score from Baseline:

319 (62.4%) vs 332 (65.6%); RR 1.05 (95% CI 0.96-1.15); p=0.29

CGI Improvement:

356 (69.7%) vs 376 (74.3%); RR 1.07 (95% CI 0.99-1.15); p=0.10

Switch Group (N=511) Vs Augment-Aripiprazole Group (N=505)

Remission:

114 (22.3%) vs 146 (28.9%); RR 1.30 (95% CI 1.05-1.60); p=0.02

$\geq 50\%$ Reduction in QIDS-C16 Score from Baseline:

319 (62.4%) vs 375 (74.3%); RR 1.19 (95% CI 1.09-1.29); p<0.001

CGI Improvement:

356 (69.7%) vs 400 (79.2%); RR 1.14 (95% CI 1.06-1.22); p<0.001

Augment-Bupropion (N=506) Vs Augment-Aripiprazole Group (N=505)

Remission:

136 (26.9%) vs 146 (28.9%); RR 1.08 (95% CI 0.88-1.31); p=0.47

$\geq 50\%$ Reduction in QIDS-C16 Score from Baseline:

332 (65.6%) vs 375 (74.3%); RR 1.13 (95% CI 1.04-1.23); p=0.003

CGI Improvement:

376 (74.3%) vs 400 (79.2%); RR 1.07 (95% CI 1.00-1.14); p=0.07

Limitations:

- All patients were from the VA Health System – external validity limited
- Open-label trial design (lack of blinding)
- Short trial duration – cannot extrapolate results or make assumptions regarding long term effects of treatment
- Trial results must be considered in relation to previous/current pharmacotherapy (primarily SSRI or SNRI)

Level of Evidence: Level I – with major limitations

Recommendation: For these reasons, I do not recommend switching therapy to bupropion in patients with major depressive disorder unresponsive to current treatment. Instead, I recommend using aripiprazole (over bupropion) to augment pharmacotherapy. However, it is important to individualize treatment (particularly in psychiatric patients) and consider drug side-effect profiles when making therapy changes.

Efficacy:

- Both augment-bupropion and augment-aripiprazole groups demonstrated significantly greater rates of remission compared to the switch group
 - There was no significant difference in the remission rates between the augment groups
- The augment-aripiprazole group demonstrated significantly greater improvement in QIDS-C16 score compared to augment-bupropion

Safety:

- Rates of serious adverse effects were similar between treatment groups
- Rates of weight gain ($\geq 7\%$), somnolence and extrapyramidal symptoms were significantly higher in the augment-aripiprazole group
 - However, rates of anxiety were lowest in this treatment group

Cost:

- The cost of using aripiprazole to augment current therapy must be balanced against the cost of using other treatment options as well as the value provided to the patient in terms of improved quality of life

Special Considerations/Populations:

- While the duration was not extensive and the patient population limits external validity the results do help provide useful information on how to guide pharmacotherapy choices in patients that are not sufficiently responsive
- Open-label trial design raises the potential for patient and investigator bias

Grade of Recommendation: **B**

“Good doctors use both individual clinical expertise and the best available external evidence, and neither alone is enough.”

-David Sackett, MD

While one’s clinical expertise increases with experience and practice, the ability to grow and maintain a working knowledge of available research and trials is severely limited by the finite hours per day, few of which can be freely dedicated to said task.

High-Powered Medicine includes concise, clinically relevant data with evidence-based recommendations from over 150 landmark trials spanning more than 30 years which can then be paired with one’s own clinical expertise to optimize pharmacotherapy in chronic disease management.